

The evolution of magma during continental rifting: new constraints from the isotopic and trace element signatures of silicic magmas from Ethiopian volcanoes

William Hutchison^{1,2}, Tamsin A. Mather¹, David M. Pyle¹, Adrian J. Boyce³, Matthew L. M. Gleeson⁴, Gezahegn Yirgu⁵, Jon D. Blundy⁶, David J. Ferguson⁷, Charlotte Vye-Brown⁸, Ian L. Millar⁹, Kenneth W.W. Sims¹⁰ and Adrian A. Finch²

¹ Department of Earth Sciences, University of Oxford, South Parks Road, Oxford OX1 3AN, UK. ² School of Earth and Environmental Sciences, University of St. Andrews, KY16 9AL, UK. ³ Scottish Universities Environmental Research Centre, Rankine Avenue, East Kilbride G75 0QF, UK. ⁴ Department of Earth Sciences, University of Cambridge, Downing Street, Cambridge CB2 3EQ, UK. ⁵ School of Earth Sciences, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia. ⁶ School of Earth Sciences, University of Bristol, Wills Memorial Building, Queens Road, Bristol BS8 1RJ, UK. ⁷ School of Earth and Environment, University of Leeds, Leeds LS2 9JT, UK. ⁸ British Geological Survey, The Lyell Centre, Research Avenue South, Edinburgh EH14 4AP, UK. ⁹ NERC Isotope Geosciences Laboratory, Keyworth, Nottingham, NG12 5GG, UK. ¹⁰ Department of Geology and Geophysics, University of Wyoming, Laramie, WY 82071, USA.

Corresponding author: William Hutchison (wh39@st-andrews.ac.uk)

Highlights:

- First ‘rift-scale’ analysis of silicic magmagenesis in Ethiopia
- Limited assimilation of high- $\delta^{18}\text{O}$ Pan-African basement in Quaternary magmas

- Fractional crystallization is variable and amplified in regions of low magma flux
- We predict ~45 Ma of magmatic intrusion has removed fusible components from crust
- Greater crustal assimilation for Oligocene magmas supports secular variations

Keywords: rift; magmatism; assimilation; peralkaline; Ethiopia; oxygen isotopes

Abstract

Magma plays a vital role in the break-up of continental lithosphere. However, significant uncertainty remains about how magma-crust interactions and melt evolution vary during the development of a rift system. Ethiopia captures the transition from continental rifting to incipient sea-floor spreading and has witnessed the eruption of large volumes of silicic volcanic rocks across the region over ~45 Ma. The petrogenesis of these silicic rocks sheds light on the role of magmatism in rift development, by providing information on crustal interactions, melt fluxes and magmatic differentiation. We report new trace element and Sr-Nd-O isotopic data for volcanic rocks, glasses and minerals along and across active segments of the Main Ethiopian (MER) and Afar Rifts. Most $\delta^{18}\text{O}$ data for mineral and glass separates from these active rift zones fall within the bounds of modelled fractional crystallization trajectories from basaltic parent magmas (i.e., 5.5–6.5 ‰) with scant evidence for assimilation of Pan-African Precambrian crustal material ($\delta^{18}\text{O}$ of 7–18 ‰). Radiogenic isotopes ($\epsilon_{\text{Nd}}=0.92\text{--}6.52$; $^{87}\text{Sr}/^{86}\text{Sr} = 0.7037\text{--}0.7072$) and incompatible trace element ratios ($\text{Rb}/\text{Nb} < 1.5$) are consistent with $\delta^{18}\text{O}$ data and emphasize limited interaction with Pan-African crust. However, there are important regional variations in melt evolution revealed by incompatible elements (e.g., Th and Zr) and peralkalinity (molar

Na₂O+K₂O/Al₂O₃). The most chemically-evolved peralkaline compositions are associated with the MER volcanoes (Aluto, Gedemsa and Kone) and an off-axis volcano of the Afar Rift (Badi). On-axis silicic volcanoes of the Afar Rift (e.g., Dabbahu) generate less-evolved melts. While at Erta Ale, the most mature rift setting, peralkaline magmas are rare. We find that melt evolution is enhanced in less mature continental rifts (where parental magmas are of transitional rather than tholeiitic composition) and regions of low magma flux (due to reduced mantle melt productivity or where crustal structure inhibits magma ascent). This has important implications for understanding the geotectonic settings that promote extreme melt evolution and, potentially, genesis of economically-valuable mineral deposits in ancient rift-settings. The limited isotopic evidence for assimilation of Pan-African crustal material in Ethiopia suggests that the pre-rift crust beneath the magmatic segments has been substantially modified by rift-related magmatism over the past ~45 Ma; consistent with geophysical observations. We argue that considerable volumes of crystal cumulate are stored beneath silicic volcanic systems (> 100 km³), and estimate that crystal cumulates fill at least 16–30 % of the volume generated by crustal extension under the axial volcanoes of the MER and Manda Hararo Rift Segment (MHRS) of Afar. At Erta Ale only ~1 % of the volume generated due to rift extension is filled by cumulates, supporting previous seismic evidence for a greater role of plate stretching in mature rifts at the onset of sea-floor spreading. We infer that ~45 Ma of magmatism has left little fusible Pan-African material to be assimilated beneath the magmatic segments and the active segments are predominantly composed of magmatic cumulates with $\delta^{18}\text{O}$ indistinguishable from mantle-derived melts. We predict that the $\delta^{18}\text{O}$ of silicic magmas should converge to mantle values as the rift continues to evolve. Although current data are limited, a comparison with ~30 Ma ignimbrites (with $\delta^{18}\text{O}$ up

64 to 8.9 ‰) supports this inference, evidencing greater crustal assimilation during initial stages of
65 rifting and at times of heightened magmatic flux.

1. Introduction

Magmatism fundamentally alters the thermal, chemical and mechanical properties of the crust and plays a key role in the break-up of continental lithosphere (Buck, 2006; Bialas et al., 2010). However, uncertainty remains about whether magmatic differentiation and crustal interactions vary spatially between different rift segments, and whether there are significant secular variations during rift evolution. Studies of the petrogenesis of rift magmas offer insights into these questions.

The petrologic diversity of volcanic rocks is generated by numerous processes. Among the most important are: interaction with the crust via partial melting or assimilation; and fractional crystallization of the parental magma (e.g., Macdonald et al., 2008; Deering et al., 2008). Partial crustal melting strongly depends on the thermal state of the crust (Dufek and Bergantz, 2005; Annen et al., 2006) and the availability of fusible crustal materials. In active rifts, the potential for partial melting will be amplified in regions of elevated temperatures and will coincide with zones of highest magmatic intrusion (Karakas and Dufek, 2015). Partial melting is also favoured in regions of fusible crust, while more refractory regions, that have already been heavily intruded, are less likely to be remelted or assimilated by later intrusions. Fractional crystallization will be amplified in rift settings where magma flux and crustal temperatures are lower, and there is an absence of fusible crust.

Geochemical techniques can discriminate between partial crustal melting and fractional crystallization. Oxygen isotopes ($\delta^{18}\text{O}$) are a powerful tool for investigating crustal interactions (provided the $\delta^{18}\text{O}$ of crust is distinct from mantle-derived rocks and cumulates), while incompatible trace elements (e.g., Ba, Sr, Th, Zr) are particularly sensitive to fractional

crystallization. Geochemical studies in active rift zones, notably Iceland, have successfully linked silicic magma petrogenesis to the thermal state of the crust (Martin and Sigmarsson, 2010). On the axis of the Icelandic Rift, where magma flux is high and the crust is hot, silicic magmas exhibit $\delta^{18}\text{O}$ evidence for assimilation of fusible hydrothermally-altered metabasaltic crust with low $\delta^{18}\text{O}$ ($< 2\text{‰}$). While in cooler off-rift settings, magmatic flux is lower, assimilation is limited (samples exhibit normal magmatic $\delta^{18}\text{O}$, 5.0–6.5‰, Eiler, 2001), and silicic melts undergo extensive fractional crystallization. In continental rift zones further complexity is expected because vestigial pre-rift continental crust may also be present.

Ethiopia exposes several stages of rift development from continental rifting in the Main Ethiopian Rift (MER) to nascent seafloor spreading in the Afar Rift (Figure 1, Hayward and Ebinger, 1996), providing a unique opportunity to study connections between magma petrogenesis and geotectonic setting. Here, geochemical data can be interpreted in the context of geophysical constraints on crustal structure and composition (Keranen et al., 2004; Bastow and Keir, 2011; Hammond et al., 2011), and magmatic intrusion volumes (Dessisa et al., 2013; Keir et al., 2015). Further, magmatism in Ethiopia has been taking place since ~45 Ma (Rooney, 2017) permitting the development of a temporal understanding of magma evolution and crustal interactions as rifting proceeds.

Previous studies in Ethiopia focused on geochemistry of mafic magmas of the MER and found evidence for spatio-temporal variations in crustal assimilation and fractionation (Rooney et al., 2007; Section 2). However, silicic volcanism is a key component of rift magmatism and a common feature across different rift zones. Although previous authors have investigated individual complexes (e.g., Gedemsa, Peccerillo et al., 2003; Dabbahu, Field et al., 2013) it is

unclear whether silicic magmagenesis varies spatially across different rift settings and whether there have been secular variations since the onset of rifting. Answering these questions has important implications for understanding ongoing rift volcanism; and the links between petrogenesis, rift setting and mineral resources. Silicic melts generated in continental rifts by protracted fractional crystallization tend to be enriched in economically-valuable elements (including, rare earth elements, REE, Zr, Nb and Ta). Identifying rift settings that favour extreme differentiation (i.e., mature versus immature continental rifts, or on- versus off-axis locations) provides valuable insights into the geotectonic settings that may host economically significant ore bodies.

In this paper we integrate new and published Sr-Nd-O isotope and trace element data from six MER and Afar Rift volcanic systems (Figure 1a, b). We evaluate the relative importance of fractional crystallization and crustal melting at each and compare this to their rift setting (crustal thickness and crustal compositions, Figure 1) and eruptive flux. We show that:

- i) despite significant variations in magma flux and crustal structure there is limited evidence for Pan-African crustal assimilation in Ethiopian Quaternary magmas
- ii) there are variations in fractional crystallization between the different volcanic systems, and melt evolution is amplified in less mature rifts with lower magma flux
- iii) the relative importance of fractional crystallization and crustal melting in the genesis of silicic magmas should vary as a continental rift develops and the pre-rift crust is modified by magmatic intrusions

2. Geological Setting

Magmatic activity in East Africa began in the Eocene. Recent reviews (Rooney, 2017) suggest multiple pulses of magmatism since ~45 Ma, with the most volumetrically-significant flood basalt and silicic eruptions taking place in the Oligocene (~33.9 to 27 Ma, Hofmann et al., 1997; Ayalew et al., 2002). Rift magmas have intruded through a continental lithosphere that comprises Precambrian schists and granitoids assembled during the Neoproterozoic Pan-African crust building event (Teklay et al., 1998). Initiation of major rift zones was diachronous: ~35 Ma in the Gulf of Aden (d'Acremont et al., 2005); ~28 Ma in the Red Sea (Wolfenden et al., 2005) and 15–18 Ma in the MER (Wolfenden et al., 2004). Each rift zone shows a comparable evolutionary history, with early deformation accommodated on border faults, and later extension and magmatic intrusions localized along 20 km wide and 60–80 km long magmatic segments (Ebinger, 2005). Geological and geophysical evidence for crustal thinning (Maguire et al., 2006; Hammond et al., 2011, Bastow and Keir, 2011), intruded magma volumes (Keranen et al., 2004; Keir et al., 2015) and rift architecture (Agostini et al., 2011) suggest rift maturity varies from intermediate-mature continental rifting in the MER to incipient seafloor spreading in Afar.

Quaternary volcanism in Ethiopia is strongly bimodal; basalts (mantle melts generated at significant depths, >80 km, and elevated temperatures, Rooney et al., 2012a; Ferguson et al., 2013a; Armitage et al., 2015) are associated with dykes and fissure eruptions, whereas rhyolites and trachytes are associated with shield-like complexes and calderas.

We focus on six volcanic systems (Figure 1a). Aluto, Gedemsa and Kone are located along the MER (Kone is in a different magmatic segment from Aluto and Gedemsa; Ebinger and Casey, 2001). In the Afar Rift, Dabbahu and Badi are located on and off the Manda Hararo Rift Segment (MHRS), respectively (Figure 1b), while the northerly Erta Ale range comprises the

153 Erta Ale Segment (EAS, Beyene and Abdelsalam, 2005). Table 1 summarizes the setting and
154 eruptive history of the volcanic systems. Based on published data, each volcanic system spans a
155 wide compositional range (45–75 wt. % SiO₂), and is represented predominantly by basalt and
156 rhyolite compositions (Figure 2a). Although rocks with intermediate silica contents are relatively
157 scarce there is a continuum of compositions (c.f. Macdonald et al., 2008) and only Kone (Figure
158 1a) completely lacks intermediate magmas (Figure 2a). Basalts from Erta Ale are notably more
159 tholeiitic than the other complexes, and maintain lower alkalinity throughout the differentiation
160 sequence.

161 Crustal thickness varies markedly between different rift zones, from ~16 km in the EAS to 20–22
162 km beneath the MHRS, and from ~40 km in the MER beneath Aluto to ~30 km beneath Kone
163 (Maguire et al., 2006, Figure 1c). The upper crust comprises vestigial Pan-African crust (Figure
164 1c, Makris and Ginzburg 1987; Mackenzie et al., 2005; Maguire et al., 2006; Hammond et al.,
165 2011) with $\delta^{18}\text{O}$ of 7–18 ‰ (Duffield et al., 1997; Ayalew et al., 2002), higher than typical
166 mantle-derived magmas (5–6.5‰, Section 5.1). Geophysical surveys suggest that the Pan-
167 African crust has been significantly modified by intrusions (particularly beneath magmatic
168 segments, Hammond et al., 2011). This is supported by geochemical studies of mafic lavas
169 which show, firstly, that crustal assimilation in Quaternary lavas is only identified in less mature,
170 more southerly, MER rift sections; secondly, that crustal assimilation is more pronounced in
171 older lava series (30 and 11–6 Ma) compared to recent samples (Rooney et al., 2007). Silicic
172 magmas, the topic of this study, have a longer residence in the crust and provide a
173 complementary and potentially more accentuated geochemical record of magma-crust interaction
174 and fractionation.

3. Methods

3.1 Analytical methods

$\delta^{18}\text{O}$ analysis of glass and mineral separates (1–2 mg) was carried out at Scottish Universities Environmental Research Centre, East Kilbride by laser fluorination following the method of Sharp (1990) modified for ClF_3 (Macaulay et al., 2000). For mineral separates, overnight prefluorination was carried out to remove adsorbed environmental water from the sample chamber and line. For glasses, which are more reactive in ClF_3 , we employed a short (90 second) room-temperature prefluorination before each analysis (Pope et al., 2013). Standards were run after each unknown and their reproducibility errors, including mass spectrometry, was typically better than ± 0.3 ‰, reported in standard notation as permil (‰) variations to V-SMOW. New analyses are compiled in Table 2. To complement the $\delta^{18}\text{O}$, a small number of samples were analysed for Sr-Nd-Pb isotopes. Detailed information on the preparation and analysis of these samples is provided in the Supplementary Information with a compilation of all whole-rock major and trace element data used here (Supplementary Data).

3.2 Thermodynamic and oxygen isotope modelling

To examine whether evolved peralkaline magmas could be generated via closed-system fractional crystallization only, we modelled potential differentiation sequences using Rhyolite-MELTS (Gualda et al., 2012). Using a primitive parental basalt composition (Table 3) and assuming isobaric fractional crystallization, we calculated the stable phase assemblage, at given pressure (P), temperature (T) and oxygen fugacity ($f\text{O}_2$), most closely matching the composition

of natural samples. We focused modelling on Aluto and Dabbahu as their sample suites have been analysed in greatest detail (Field et al., 2013; Gleeson et al., 2017). A range of parameters was explored (Table 3), and a minimization routine used to identify the best-fit P, T and fO_2 conditions matching whole-rock data (Gleeson et al., 2017). While there are well-known limitations applying Rhyolite-MELTS to peralkaline systems (discussed by Rooney et al., 2012b; Gleeson et al., 2017), models provide a reasonable fit to the compositional data and are sufficient to gain first-order understanding of the liquid lines of descent and crystallization sequence required to generate silicic peralkaline melts.

Few studies have investigated the variation of $\delta^{18}O$ in peralkaline magma. We model the expected changes in $\delta^{18}O_{\text{melt}}$ during closed-system fractional crystallization using the approach of Bindeman et al. (2004). Taking the step-wise crystallizing assemblage, temperature and melt composition from the best-fitting Rhyolite-MELTS model, we calculate $\delta^{18}O_{\text{cumulate}}$ and subtract this from the $\delta^{18}O_{\text{melt}}$ value. We treated the melt as a mixture of CIPW normative minerals, and calculate the temperature and melt composition-dependent mineral-melt fractionations ($\Delta_n^{i+1}(T)$) at each step (i+1) and for each crystallizing mineral (n). This forward-step mass balance model (detailed in the Supplementary Information) follows equations from Bucholz et al. (2017). We determine $\delta^{18}O_{\text{melt}}$ at each stage of peralkaline melt genesis from primitive rift-related basalts and predict the $\delta^{18}O$ trajectory that may plausibly represent the products of closed-system fractional crystallization. Samples that fall off the modelled $\delta^{18}O_{\text{melt}}$ fractionation trajectory have likely assimilated local crust (see Section 5.1).

4. Results

4.1 Major element trends and Rhyolite-MELTS models

Silicic rocks from Ethiopia are mainly peralkaline (i.e., molar $\text{Na}_2\text{O} + \text{K}_2\text{O} / \text{Al}_2\text{O}_3$, NK/A , >1 ; Figure 2b). The most peralkaline samples ($\text{NK/A} > 1.6$) are associated with the MHRS off-axis volcano Badi and MER volcanoes Kone, Gedemsa and Aluto. Erta Ale samples are only mildly peralkaline, while peraluminous rocks (i.e., molar $\text{Al}_2\text{O}_3 / \text{CaO} + \text{K}_2\text{O} + \text{Na}_2\text{O} > 1$) are found at Gedemsa (Figure 2b). The volcanic systems are more clearly distinguished using the peralkaline classification diagram of Macdonald et al. (1974) that shows all complexes, except Erta Ale, are dominated by pantellerites, with lesser comendites (Figure 2c). Kone, Gedemsa, Aluto and Badi erupt the most evolved pantelleritic melts (Figure 2c).

Major element trends overlap (Figure 3), suggesting a similar pattern of crystallization and melt evolution at each system. The most obvious major element differences are observed in samples with >70 wt. % SiO_2 . Rhyolites show considerable scatter in FeO_t values, reflecting varying degrees of fayalite, alkali pyroxene, aenigmatite and Fe-Ti oxide removal or accumulation in the final stages of melt evolution. Rhyolites with anomalously low Na_2O (Gedemsa and Kone samples, Figure 3) tend to have high loss on ignition, perhaps reflecting post-emplacement alteration (Peccerillo et al., 2003). There is considerable scatter in Al_2O_3 above 70 wt. % SiO_2 suggesting that feldspar fractionation is highly variable, while the Al_2O_3 minima suggest that more extensive feldspar fractionation occurred at Kone, Gedemsa, Aluto and Badi compared to Dabbahu and Erta Ale (Figure 3). P_2O_5 for most suites falls on a non-linear trend with an inflection at ~ 55 wt. % SiO_2 that reflects stabilization of apatite (Rooney et al., 2012b; Field et al., 2013). The behaviour of P_2O_5 suggests that fractional crystallization is the main process generating the magmas (c.f. Lee and Bachmann, 2014), although a few enclaves from Gedemsa

(Peccerillo et al., 2003) and basaltic trachyandesites from Dabbahu (Field et al., 2011) fall along trends consistent with magma-mixing (Figure 3).

Rhyolite-MELTS fractional crystallization models for Aluto and Dabbahu reproduce reasonably well the trends observed in whole-rock data (Figure 3). In both cases the best-fit models were able to generate pantellerite melts from the most primitive mafic samples at low pressures (150 MPa), low initial H₂O concentrations (~0.5 wt. %) and relatively low fO₂ (QFM; Table 3).

Rhyolite-MELTS modelling is consistent with pantellerites being produced by protracted fractional crystallization (>80 %) of primitive rift-related basalts (Gleeson et al., 2017).

Discrepancies between Rhyolite-MELTS models and whole-rock data are generally restricted to the final stages of crystallization (>65 wt. % SiO₂), as explored by Rooney et al. (2012) and Gleeson et al. (2017). In short, the Rhyolite-MELTS apatite solubility model overpredicts P₂O₅ for peralkaline magmas throughout fractionation (Rooney et al., 2012b). CaO is also overpredicted, linked to inaccuracies in the stabilization of apatite (Rooney et al., 2012b). FeO_t concentrations for Rhyolite-MELTS models are 1–6 wt. % lower than natural sample values at high SiO₂ (>65 wt. %), reflecting the limited constraints on aenigmatite stability. These inaccuracies tend to be associated with volumetrically minor phases (e.g., aenigmatite and apatite: <5%, Field et al., 2013; Gleeson et al., 2017), and best-fit liquid lines of descent (Figure 3) capture the trend of natural samples over most of their differentiation.

259 **4.2 Trace element trends**

260 Each volcanic system exhibits extreme compatible (e.g., Sr) and incompatible (e.g., REE, Nb)
261 trace element variation, consistent with protracted fractional crystallization (Figure 4). Zircon
262 solubility in peralkaline melts is significantly higher than sub-alkaline melts (Watson, 1979),
263 consequently Zr remains almost perfectly incompatible during differentiation in peralkaline
264 rhyolites (Field et al., 2012a). Thus, we can use Zr as a marker of degree of magmatic evolution.
265 In each panel of Figure 4 there is a near-continuous trend from primitive mafic lavas (lowest Zr,
266 ~100 ppm) to silicic rocks (highest Zr, up to ~3000 ppm). Sr is compatible in both plagioclase
267 and alkali feldspar and there is a clear distinction between the relatively high Sr contents of Erta
268 Ale silicic rocks and the low concentrations elsewhere (Figure 4). Low Ba concentrations (<500
269 ppm) in rhyolites reflect alkali feldspar removal and are observed at all volcanic systems except
270 Erta Ale, which lacks alkali feldspar (Figure 4; Bizouard et al., 1980). Silicic samples from Aluto
271 have relatively high Ba (180–470 ppm), Gedemsa and Kone are similar in terms of their minima
272 (~25 ppm) and pattern, while the lowest Ba values (~3 ppm) are observed at Dabbahu and Badi.
273 Several silicic samples from Gedemsa with elevated Ba have previously been linked to alkali
274 feldspar accumulation (Peccerillo et al., 2003).

275 The volcanic systems are characterized by subtly different but generally constant incompatible
276 trace element ratios, e.g., Nb/Zr and Rb/Zr (Figure 4). They also exhibit clear distinctions in
277 maximum incompatible element enrichment (Figure 4). The greatest incompatible element
278 concentrations are observed at the MER volcanoes Kone (~2400 ppm Zr) and Gedemsa (~2300
279 ppm Zr), and the MHRS off-axis volcano Badi shows the largest range with maximum ~3000

ppm Zr. Zr contents of Erta Ale samples reach only ~750 ppm (Figure 4). Zr contents of Dabbahu samples reach 1300 ppm and divide into a high and low Nb series at ~700 ppm Zr. Dabbahu samples displaced to lower Nb values are comendites, possibly reflecting a greater role of ilmenite (which accommodates Nb) in their formation. However, we acknowledge that the P-T-fO₂ and compositional conditions that govern Fe-Ti oxide stability in peralkaline melts remain poorly understood (Marshall et al., 2009) and it is unclear what magmatic processes might favour ilmenite crystallization in these samples.

Y is positively correlated with Zr for all complexes except Badi, where several samples fall off the linear array (Figure 4), perhaps due to amphibole and/or apatite fractionation (c.f. White et al., 2009). Th is highly incompatible in peralkaline magmas (Martin and Sigmarsson, 2010) and samples from Dabbahu, Badi, Erta Ale and Aluto define near constant Th/Zr ratios. Gedemsa samples are scattered, with elevated Th/Zr and Pb/Zr across the compositional range, likely reflecting variability in the parental basalts (c.f. Giordano et al., 2014). At the rift-scale, minor differences in incompatible element ratios between the MER and Afar Rift may represent subtle differences in parental magmas between regions. Ratios sensitive to crustal assimilation of Pan-African basement (Rb/Nb, Figure 4) are low for Quaternary volcanic systems but clearly elevated for ~30 Ma ignimbrites (Ayalew et al., 2002).

4.3 Isotopic constraints

$\delta^{18}\text{O}$ results are presented in Figure 5a. We consider all $\delta^{18}\text{O}_{\text{glass}}$ values to be representative of melt composition and have used experimentally determined fractionation factors of Appora et al.

(2003) to estimate and plot $\delta^{18}\text{O}_{\text{melt}}$ from the measured $\delta^{18}\text{O}_{\text{mineral}}$ (olivine, quartz and feldspars).
Erta Ale values after Barrat et al. (1998) are derived from whole-rock $\delta^{18}\text{O}$. As their crystal
content is low (~20 % by volume, Field et al., 2012b) it is reasonable to assume that $\delta^{18}\text{O}_{\text{whole-rock}}$
provides a close approximation to $\delta^{18}\text{O}_{\text{melt}}$.

The vast majority of $\delta^{18}\text{O}$ values fall between 5 and 6.5 ‰, and lie within the range predicted by
fractional crystallization of primitive rift-related magmas (shaded area, Figure 5a, detailed
discussion in Section 5.1). High $\delta^{18}\text{O}_{\text{melt}}$ (>7.0 ‰) is observed in only three Quaternary samples:
a Kone pre-caldera silicic sample; a Gedemsa alkali feldspar separated from a pre-caldera
rhyolite and a Dabbahu comendite (Table 2). The most evolved samples from Erta Ale divide
into two populations, with normal (5.5–6.4 ‰) and low $\delta^{18}\text{O}$ (down to 5.2 ‰, Barrat et al.,
1998). New and published $\delta^{18}\text{O}_{\text{olivine}}$ data are within error of typical upper mantle values (5.2 ± 0.2
‰, Eiler, 2001). Further, $\delta^{18}\text{O}$ values for ~30 Ma ignimbrites (grey crosses, after Ayalew et al.,
2002) show both normal and high $\delta^{18}\text{O}_{\text{melt}}$ groupings.

New Sr- and Nd-isotope results from the MHRS, Dabbahu and Badi are combined with previous
analyses in Figure 5b. Sr-Nd isotopes are available for all systems except Aluto. ϵ_{Nd} values range
from 6.52 to 0.92 with generally close agreement between the Nd-isotopic ratios for silicic and
mafic samples from the same complex. Most samples fall within the Ethiopian mantle array
(dashed black ellipse) although a number of silicic samples are elevated to more radiogenic
 $^{87}\text{Sr}/^{86}\text{Sr}$ values (up to 0.7072). Finally, Sr- and Nd-isotopic ratios for ~30 Ma basalts and
ignimbrites from across Ethiopia show greater scatter and overlap with Pan-African crustal
samples (Figure 5b).

5. Discussion

5.1 Fingerprinting crustal interactions: $\delta^{18}\text{O}$ constraints

Oxygen isotopes can provide valuable insights into magma-crust interactions. However, it is instructive to first consider how fractional crystallization affects the $\delta^{18}\text{O}$ of peralkaline magmas. Most $\delta^{18}\text{O}$ measurements from natural volcanic series show typical mantle-derived basalts have $\delta^{18}\text{O}$ of 5.2–5.8 ‰ (Eiler, 2001) with values increasing a little during fractional crystallization (Harris et al., 2000). We model the $\delta^{18}\text{O}_{\text{melt}}$ trajectory for peralkaline rhyolites in Ethiopia from Aluto and Dabbahu (Section 3.2) and find that evolved rhyolites should show a modest $\delta^{18}\text{O}_{\text{melt}}$ increase of ~0.6 ‰ during fractional crystallization from basalt (Figure 5a). The minor $\delta^{18}\text{O}_{\text{melt}}$ increase is explained because Rhyolite-MELTS predicts that the bulk (~80 %) of the fractionating assemblage comprises mineral phases (olivine, clinopyroxene and plagioclase) that have relatively small fractionation factors at temperatures of 1250–750 °C (Eiler, 2001).

The modelled $\delta^{18}\text{O}_{\text{melt}}$ trajectory is shaded in Figure 5a. We interpret samples that fall outside this range as having interacted with high or low $\delta^{18}\text{O}$ sources. We emphasize that the $\delta^{18}\text{O}$ of magmatic residue or cumulate phases overlaps with normal magmatic $\delta^{18}\text{O}$ and so interactions with these cannot be detected by $\delta^{18}\text{O}$ analyses (see Section 5.4). High $\delta^{18}\text{O}$ samples have assimilated Pan-African crustal rocks (Figure 1c) with $\delta^{18}\text{O}$ of 7–18 ‰ (Duffield et al., 1997; Ayalew et al., 2002) while low $\delta^{18}\text{O}$ samples have likely assimilated shallow hydrothermally-altered crustal rocks. We rule out a sub-solidus alteration or weathering cause for low $\delta^{18}\text{O}$ lavas because sampling targeted exceptionally fresh material without visible surface alteration (Barrat et al., 1998). Further, while in principle low $\delta^{18}\text{O}$ silicic magmas could be generated by fractional

crystallization of low $\delta^{18}\text{O}$ mafic magmas, none of the basaltic samples or olivine separates (Table 2, Figure 5a) lie significantly outside typical upper mantle range (Section 4.2).

High-temperature hydrothermal circulation of low- $\delta^{18}\text{O}$ meteoric fluids at active volcanic systems causes alteration and generates low- $\delta^{18}\text{O}$ volcanic crust (Norton and Taylor, 1979). Geothermal activity is reported at all volcanic systems in the study, geothermal fluids and groundwater have $\delta^{18}\text{O}$ between -1 and -5 ‰ (Duffield et al., 1997; Darling et al., 1996), and at Aluto deep drilling has identified high temperatures (300 – 400 °C) and extensive hydrothermally-altered volcanic deposits (Teklemariam et al., 1996). To date, there are no $\delta^{18}\text{O}$ measurements of the hydrothermally-altered facies underlying Ethiopian volcanoes, however, a useful comparison can be made with well-studied geothermal systems in Iceland such as Krafla. Here, Pope et al. (2013) showed that shallow altered volcanic crust has values of -6 ‰, intermediate between unaltered basalts (~ 5.5 ‰) and geothermal fluids (-12 ‰). Unaltered Ethiopian basalts are also ~ 5.5 ‰ (Figure 5) and so assuming comparable levels of fluid-rock interaction to Iceland we estimate that Ethiopian hydrothermally-altered crust has $\delta^{18}\text{O}$ of 2 ‰ (we averaged all $\delta^{18}\text{O}$ values from Duffield et al., 1997 and Darling et al., 1996, above, to make a conservative estimate of -1.5 ‰ for geothermal fluids).

In Figure 5a, the horizontal lines above and below the grey shaded fractional crystallization $\delta^{18}\text{O}_{\text{melt}}$ trajectory indicate the percentage of melt sourced from crustal materials assuming end-member values of 2 ‰ (hydrothermally-altered volcanic crust) and 12 ‰ (Pan-African crust). Until detailed $\delta^{18}\text{O}$ studies of local crust are undertaken at each system (i.e., from xenoliths) our estimates of melt mass percentage serve guidelines rather than precise values.

5.2 Genesis of silicic melts in Quaternary magmatic segments

The $\delta^{18}\text{O}$ of Quaternary volcanic products from across Ethiopia generally fall between 5.0 and 6.5 ‰ and show a close correspondence with our modelled $\delta^{18}\text{O}_{\text{melt}}$ fractional crystallization trajectory (Figure 5a). The pantellerites show no oxygen-isotopic evidence for crustal assimilation, strongly suggesting that the most evolved silicic rocks of Ethiopia are not derived from melting of high- $\delta^{18}\text{O}$ Pan-African crust. This conclusion agrees with trace element evidence, for example, large ion lithophile element and high field strength element ratios (LILE/HFSE, e.g. Rb/Nb <1.5) which show only limited overlap with Pan-African (Figure 4, Peccerillo et al., 2003, 2007).

Subtle crustal assimilation signatures are detected in a limited number of Quaternary samples (Figure 5a), and Sr-Nd isotopes provide an additional test (Figure 5b). Sr concentrations in the silicic samples are very low (Figure 4) making Sr-isotopes very sensitive to crustal assimilation. In Figure 5b a simple binary mixing model is shown for a typical uncontaminated mantle-derived melt (white star) and a Pan-African lithosphere end-member (grey star, from Rooney et al., 2012c). The mixing curves demonstrate that for a typical pantelleritic melt (starting with 5 ppm Sr and 200 ppm Nd) addition of 5–10 % Pan-African material can displace $^{87}\text{Sr}/^{86}\text{Sr}$ to high crustal values without significantly altering ϵ_{Nd} . Thus, elevated $^{87}\text{Sr}/^{86}\text{Sr}$ observed in various silicic samples in Figure 5b are explained by relatively low levels of crustal assimilation (<10 %). This agrees with previous radiogenic isotope studies in the MER (e.g., Peccerillo et al., 2003; Giordano et al., 2014) that identified elevated $^{87}\text{Sr}/^{86}\text{Sr}$ in feldspar phenocrysts (up to ~0.7056) as well as whole-rock samples.

Together, the major and trace elements as well as the Sr-Nd-O isotopes and Rhyolite-MELTS models are consistent with fractional crystallization from basalts to rhyolites with very minimal crustal assimilation to explain the origins of Quaternary peralkaline magmas across Ethiopia. These observations agree with petrogenetic studies from other regions of continental extension, e.g., Taupo Volcanic Zone, where fractional crystallization is the primary mechanism of silicic magmagenesis (e.g., Graham et al. 1995; Deering et al., 2008).

The three high- $\delta^{18}\text{O}$ samples (Table 1) relate to pre-caldera phases of activity (samples GD22 and 2-11 from Gedemsa and Kone, respectively) and a period of comendite melt production following a hiatus in silicic activity (sample 011, Dabbahu). $\delta^{18}\text{O}$ data from Aluto do not follow a similar temporal trend as both pre-caldera (6.5 ‰) and early post-caldera comenditic samples (6.1 ‰) are very close to the recent post-caldera products (6.2–6.4 ‰). Detailed discussion on the origins of these high- $\delta^{18}\text{O}$ samples is beyond the scope of this study, although future isotopic investigations of glass and minerals from these outliers is desirable.

Although the key finding from our $\delta^{18}\text{O}$ analysis is that assimilation of Pan-African crustal materials is limited, our analyses also reveal that evidence for crustal assimilation is entirely absent in some systems. For example, no high- $\delta^{18}\text{O}$ lavas were found at Erta Ale, Aluto, or Badi (Figure 5a). Erta Ale is notable because it is in the most mature rift setting. Although magmatic flux and partial crustal melting are expected to be greatest here (Section 1), the absence of a high- $\delta^{18}\text{O}$ signal strongly suggests that earlier intrusions and crustal modification have effectively removed any fusible Pan-African crust. This hypothesis is consistent with geophysical evidence and is developed with petrological models for crustal modification in Section 5.4. An additional feature of the $\delta^{18}\text{O}$ data is that, despite large geothermal systems being

present at each volcanic system studied, only at Erta Ale are low $\delta^{18}\text{O}$ silicic samples identified (trachytes, Figure 5a). In this case we suspect that hydrothermally-altered crust in the EAS (Figure 1a) might be more extensive and/or altered based on seismic interpretations indicating thicker rift fill (Figure 1c, Hammond et al., 2011) and the more mature rift morphology and pervasive faulting facilitating greater meteoric water ingress.

5.3 Geotectonic controls on melt evolution

Peralkaline magmas are a ubiquitous feature of silicic volcanism in Ethiopia. However, we identified clear variations in melt evolution between the volcanic systems. The most evolved peralkaline melts from Kone, Gedemsa, Aluto and Badi are similar in terms of their exceptional peralkalinity ($\text{NK/A} > 1.6$, Figure 2b) and extreme incompatible element enrichment (e.g., Zr and Th, Figure 4). By comparison, pantellerites from Dabbahu volcano are less peralkaline ($\text{NK/A} < 1.6$), with lower incompatible element concentrations, while at Erta Ale, silicic peralkaline melts are rare.

These observations beg the question: what drives extreme fractional crystallization, and what rift settings amplify these processes? In general, conditions that promote peralkaline rhyolite genesis are: (1) fractional crystallization of a transitional basalt, with an alkali content between those of tholeiitic and alkali basalts (Barberi et al., 1975); (2) low oxygen fugacity ($\text{NNO}-1$ or below; Scaillet and Macdonald 2001); (3) retention of a magmatic volatile phase which extends the crystallization interval to low pressures (≤ 150 MPa) and temperatures ($\leq 750^\circ\text{C}$) (Macdonald, 1987; Macdonald et al., 2014).

The flux of magma into peralkaline volcanic systems also plays a key role promoting geochemical variations (Macdonald, 2012). At high magma fluxes, the elevated heat and mass input promotes homogenization, reducing the likelihood of reaching the extremes of fractional crystallization (Macdonald et al., 2014). Magmatic fluxes are difficult to quantify directly and so we calculate eruptive fluxes for each volcanic system, and make the reasonable assumption that these are proportional to magmatic flux. Erupted volumes are calculated from the volume of the edifice and caldera (where present) following methods of Hutchison et al. (2016a), and fluxes are estimated using available age constraints (summarized in Table 4). Clearly, flux estimates do not consider dykes and unerupted magmas, and so a necessary assumption is that intrusive:extrusive ratios are comparable for each volcanic setting. Further, due to the paucity of detailed mapping and geochemistry on individual silicic centres along the EAS we did not estimate their volumes. Instead, we treat the segment as a single volcanic system; this is justified because the Erta Ale range has smooth along-axis topography (analogous to a single shield volcano) and the eruptive centres show clear evidence for magmatic interconnectivity (Pagli et al., 2012; Xu et al., 2017).

Eruptive volumes and fluxes are plotted in Figure 6a-b and reflect variations in mantle melt productivity as well as crustal melt storage. The greatest eruptive volumes ($\sim 400 \text{ km}^3$) are associated with Erta Ale. For the MHRS, although Dessisa et al. (2013) and Medynski et al. (2015) have reported magnetotelluric and geochemical evidence for an off-axis mantle magma reservoir developing beneath Badi over the last $\sim 30 \text{ ka}$, our volume estimates suggest that over longer ($>100 \text{ ka}$) timescales the eruptive fluxes have been greater at on-axis Dabbahu, compared to off-axis Badi. In the MER, eruptive volumes increase southwards with the greatest fluxes associated with Aluto (Figure 6b, Table 4). Although one might expect magmatic fluxes to increase northwards along the MER, and correlate with increasing rift maturity, seismic

tomography (Gallacher et al., 2016) reveals that the lowest velocities (i.e. greatest mantle melt production) are associated with the central MER (beneath the Aluto-Gedemsa segment) rather than Kone and more northern MER segments. Comparing eruptive volumes and fluxes with geochemical indicators of melt evolution (e.g., Zr concentration, Figure 6c) we find that volcanic systems with the lowest eruptive fluxes (i.e., Badi, Kone and Gedemsa, $0.05\text{--}0.2\text{ km}^3\text{ ka}^{-1}$) have the most evolved magmas.

Rift maturity is one of the main controls on melt evolution and this is demonstrated in nascent seafloor spreading settings like Erta Ale. Here, the lithosphere is extensively thinned and enhanced decompression melting (Bastow and Keir, 2011) generates parental basalts that are tholeiitic (Figure 2a) and therefore unable to generate evolved peralkaline rhyolites (White et al., 2009). In contrast, parental melts in the MER and MHRS of Afar are transitional basalts (Figures 2a, 4) and since Rhyolite-MELTS models (Table 3) suggest similar magma storage conditions we look to other explanations for differences in peralkalinity between these systems.

Our analysis of eruptive volumes strongly suggests that local magmatic flux is also an important control on peralkalinity (Figure 6c). Hence the more chemically-evolved nature of the MER volcanoes and off-axis Badi compared to those of on-axis Dabbahu results from the higher magma flux on the MHRS axis. As outlined above, variations in local magma supply reflect spatial variations in mantle magma productivity and/or variations in crustal melt storage. In the second case it is well known that variations in rift structures and stress field play an important role in magma ascent and storage (e.g., Maccaferri et al., 2014). Protracted magma storage will

promote fractionation and melt evolution, but will also favour cooling and degassing making the magma less likely to erupt.

In the MHRS, mantle melt production does not appear to be enhanced beneath the rift-axis (Dessisa et al., 2013) and so we suggest that at Badi the magma supply is lower because off-axis magma plumbing systems are less well-developed (i.e., crustal structures and/or stress field inhibit magma ascent). Thus off-axis melts at Badi are more likely to stall throughout the crust, decreasing melt supply to peralkaline magma reservoirs, which ultimately reduces homogenization and promotes more extreme fractionation. In the MER, regions of thicker crust do not amplify peralkalinity (e.g., Aluto, Figure 6c) and so we suggest that local magma supply in the MER indicates variations in mantle melt productivity. Thus melt production is greatest in the central MER beneath the Aluto-Gedemsa segment (in agreement with seismic tomography, Gallacher et al., 2016) and this reduces melt evolution compared to Kone. In summary, settings where the parental basalts are transitional in composition, and where local magmatic and thermal fluxes are low, will promote extreme melt evolution (Figure 6d).

5.4 Crustal modification

A corollary of our conclusion that fractional crystallization is the main driver of silicic magmagenesis is that large volumes of crystal cumulate are generated and stored within the crust. Estimates of cumulate volumes can be made from incompatible element contents of whole-rock samples and Rhyolite-MELTS models. Since Zr is completely incompatible (Field et al., 2012a) we use it as a proxy for melt fraction ($F = Zr_{parent} / Zr_{sample}$), assuming the most primitive

basalt at each volcano represents parental magma. Using Rhyolite-MELTS we predict cumulate phase proportions at each fractionation and temperature step, and so by using the Zr constrained melt fraction (F) it is possible to place all samples (from basalt to peralkaline rhyolite) on the Rhyolite-MELTS liquid line of descent and evaluate the crystallizing minerals and their volumes. From eruptive volume estimates (Section 5.3), and assuming sample suites from each volcano are representative, we estimate cumulate volumes and fluxes (Figure 6a-b, Table 4).

Our calculations demonstrate that the volcanic systems generate large cumulate bodies (100–450 km³, Figure 6a) and that the silicic complexes of the MER and MHRS will have cumulate volumes 3–5 times greater than the erupted volume (Table 4). At Erta Ale, where volcanic products are much less evolved and dominated by mafic samples (Figure 2a), we find that more material is erupted than forms cumulate (i.e., cumulate volume:erupted volume is 0.5, Table 4).

Magmatic cumulates must be accommodated in the extending crust. For each volcanic system we have calculated the space created by rift extension (using the crustal thickness, extension rate and the along-axis length of the system) and have compared this to the volume of cumulates needed to balance the erupted volumes. The comparison is shown in the final column of Table 4 (ratio of cumulate volume to total space created by extension). For the axial volcanoes of the MER and MHRS of Afar we calculate that 16–30 % of the volume generated by crustal extension beneath a silicic complex would be filled by magmatic cumulates. This is a considerable volume and we stress that this is a minimum because the volume fluxes do not account for unerupted (intrusive) melts (Section 5.3). Recent mafic dyking sequences in the MHRS (Ferguson et al., 2010), suggest intrusive:extrusive ratios of between 5:1 and 10:1 and so dykes and sills will also play a major role in crustal modification at the magmatic segments.

Our petrological estimates of cumulate volumes have clear geophysical consequences and it is important to underscore that there is abundant seismic evidence for large volumes of mid-crustal intrusions beneath the magmatic segments of the MER and Afar (Keranen et al., 2004; Mackenzie et al., 2005). Gravity surveys require high-density bodies ($3000\text{--}3100\text{ kg m}^{-3}$, Markris and Ginzburg, 1987; Cornwell et al, 2006; Lewi et al., 2016) to be present beneath the magmatic segments as well as the silicic complexes (including Gedemsa, Mahatsente et al., 1999). In fact, the bulk densities we calculated with Rhyolite-MELTS for cumulates ($\sim 3050\text{ kg m}^{-3}$, Table 3) are remarkably similar to the gravity-derived densities.

Geophysical and petrological evidence reinforce the notion that large volumes of the Ethiopian crust beneath the magmatic rift segments comprise magmatic intrusions and cumulates. An interesting observation from the cumulate volumes (Table 4) is that at Erta Ale only $\sim 1\%$ of the volume generated due to rift extension is filled by cumulates. This implies that extension in the EAS must be accommodated by greater levels of dyking and/or plate stretching than in the MER/MHRS. Although our petrological observations cannot discriminate between these two alternatives, seismic evidence (Bastow and Keir, 2011) clearly supports the latter revealing that the plate is abruptly thinned beneath the EAS (Figure 1c). Thus, both rift structure (from seismic) and cumulate volumes (from magma chemistry) can be used to infer a greater role of plate stretching in mature rifts at the onset of sea-floor spreading.

Finally, our interpretation that the Ethiopian crust has been substantially modified by magmatic intrusions explains why $\delta^{18}\text{O}$ evidence for Pan-African crustal assimilation is so limited (Section 5.2). $\sim 45\text{ Ma}$ of magmatism in the region has likely removed or assimilated most fusible crustal contaminants, and the active segments are predominantly composed of magmatic cumulates that

have $\delta^{18}\text{O}$ indistinguishable from the more recent mantle-derived melts, except in rare cases where hydrothermal alteration has subsequently lowered $\delta^{18}\text{O}$. We suggest that $\delta^{18}\text{O}$ of silicic magmas should converge to mantle values during rift evolution. Such a secular trend in crustal contamination has been observed in MER mafic samples (Rooney et al., 2007) and intriguingly the limited $\delta^{18}\text{O}$ data from ignimbrites associated with flood basalt volcanism at ~30 Ma (Ayalew et al., 2002) include some examples with exceptionally high $\delta^{18}\text{O}$ (Figure 5a) and low ϵ_{Nd} (Figure 5b), consistent with greater availability of crustal contaminants in earlier phases of rifting.

6. Conclusions

Silicic volcanic rocks in Ethiopia provide important insights into how magmas evolve and interact with the crust during continental break-up. Magmas of Quaternary age show little geochemical evidence for interaction with Pan-African crust. They generally have $\delta^{18}\text{O}$ of 5.5–6.5 ‰, independent of variations in crustal thickness (15–40 km), extension rates (5–15 mm a⁻¹) and eruptive fluxes (0.05–0.4 km³ ka⁻¹). Sr-Nd isotopes and trace element systematics confirm that geochemical diversity of these young silicic magmas primarily reflects variations in extent of fractional crystallization.

Peralkaline magmas associated with ancient rift zones have generated many of the world's largest REE deposits and a major challenge for exploration geologists is to identify the geotectonic settings that host economically significant ore bodies (Goodenough et al., 2016). Our analyses suggest that the two most important controls on melt evolution, and hence REE

enrichment, are rift maturity and local magma supply (Section 5.3). Thus intermediate-mature continental rifts (i.e., MER and MHRS of Afar) where parental magmas are transitional rather than tholeiitic, and where local magma supply is reduced (due to lower mantle melt productivity, e.g., Kone, and/or crustal structures or stress field inhibiting magma ascent, e.g., off-axis Badi) offer the best settings for differentiation and melt peralkalinity. Early phases of rift development might also be sufficient for generating evolved peralkaline melts, however, caution is required because in Ethiopia the high magmatic fluxes associated with ~30 Ma flood basalts and ignimbrites likely did not allow silicic melts to evolve to the same extent as many Quaternary settings (Figure 4). Our study underscores that ancient rift zones that have reached and preserved an intermediate-mature phase of development, ideally with off-axis volcanism, should provide excellent prospects for major REE deposits.

Acknowledgments, Samples, and Data

This work was supported by the RiftVolc project (NE/L013932/1, Rift volcanism: past, present and future) and a NERC studentship (NE/J5000045/1) awarded to W.H. W.H. also received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No. 689909. C. V-B. and I. M. acknowledge a BGS-NIGL Programme Project (40227) for Sr-Nd-Pb isotope analysis of the four samples from Dabbahu. C.V-B. publishes with permission of the Executive Director of the British Geological Survey (NERC). Oxygen isotope analysis at SUERC was supported by a NERC Stable Isotope Facility grant (IP-1367-0513). Technical assistance at SUERC was provided by A. MacDonald and T. Donnelly. J. Blusztajn assisted with the isotopic measurements at Woods Hole Oceanographic Institution. We thank L.

581 Field, S. Medynski and M. Rampey for providing samples and A. Peccerillo for providing
582 clarification on previous geochemical data. J. Biggs provided valuable comments on an earlier
583 version of the manuscript. We also thank Tyrone Rooney and Özge Karakas for thorough,
584 constructive reviews and Mike Bickle for editorial handling.

References

- Agostini, A., Bonini, M., Corti, G., Sani, F., Mazzarini, F., 2011. Fault architecture in the Main Ethiopian Rift and comparison with experimental models: Implications for rift evolution and Nubia–Somalia kinematics. *Earth Planet. Sci. Lett.* 301, 479–492.
doi:10.1016/j.epsl.2010.11.024
- Annen, C., Blundy, J.D., Sparks, R.S.J., 2006. The genesis of intermediate and silicic magmas in deep crustal hot zones. *J. Petrol.* 47, 505–539. doi:10.1093/petrology/egi084
- Appora, I., Eiler, J.M., Matthews, A., Stolper, E.M., 2003. Experimental determination of oxygen isotope fractionations between CO₂ vapor and soda-melilite melt. *Geochim. Cosmochim. Acta* 67, 459–471. doi:10.1016/S0016-7037(02)01090-6
- Armitage, J.J., Ferguson, D.J., Goes, S., Hammond, J.O.S., Calais, E., Rychert, C.A., Harmon, N., 2015. Upper mantle temperature and the onset of extension and break-up in Afar, Africa. *Earth Planet. Sci. Lett.* 418, 78–90. doi:10.1016/j.epsl.2015.02.039
- Ayalew, D., Barbey, P., Marty, B., Reisberg, L., Yirgu, G., Pik, R., 2002. Source, genesis, and timing of giant ignimbrite deposits associated with Ethiopian continental flood basalts. *Geochim. Cosmochim. Acta* 66, 1429–1448. doi:10.1016/S0016-7037(01)00834-1
- Ayalew, D., Yirgu, G., 2003. Crustal contribution to the genesis of Ethiopian plateau rhyolitic ignimbrites: basalt and rhyolite geochemical provinciality. *J. Geol. Soc. London.* 160, 47–56.
doi:10.1144/0016-764901-169

604 Ayalew, D., Jung, S., Romer, R.L., Kersten, F., Pfänder, J.A., Garbe-Schönberg, D., 2016.
 605 Petrogenesis and origin of modern Ethiopian rift basalts: Constraints from isotope and trace
 606 element geochemistry. *Lithos* 258–259, 1–14. doi:10.1016/j.lithos.2016.04.001

607 Barberi, F., Ferrara, G., Santacroce, R., Treuil, M., Varet, J., 1975. A transitional basalt-
 608 pantellerite sequence of fractional crystallization, the Boina Centre (Afar Rift, Ethiopia). *J.*
 609 *Petrol.* 16, 22–56. doi:10.1093/petrology/16.1.22

610 Barberi, F., Tazieff, H., Varet, J., 1972. Volcanism in the Afar Depression: Its Tectonic and
 611 Magmatic Significance. *Dev. Geotecton.* 7, 19–29. doi:10.1016/B978-0-444-41087-0.50007-5

612 Barberi, F., Varet, J., 1970. The Erta Ale volcanic range (Danakil Depression, northern Afar,
 613 Ethiopia). *Bull. Volcanol.* 34, 848–917. doi:10.1007/BF02596805

614 Barrat, J.A., Fourcade, S., Jahn, B.M., Cheminée, J.L., Capdevila, R., 1998. Isotope (Sr, Nd, Pb,
 615 O) and trace-element geochemistry of volcanics from the Erta' Ale range (Ethiopia). *J. Volcanol.*
 616 *Geotherm. Res.* 80, 85–100. doi:10.1016/S0377-0273(97)00016-4

617 Barrat, J.A., Joron, J.L., Taylor, R.N., Fourcade, S., Nesbitt, R.W., Jahn, B.M., 2003.
 618 Geochemistry of basalts from Manda Hararo, Ethiopia: LREE-depleted basalts in Central Afar.
 619 *Lithos* 69, 1–13. doi:10.1016/S0024-4937(03)00044-6

620 Bastow, I.D., Keir, D., 2011. The protracted development of the continent–ocean transition in
 621 Afar. *Nat. Geosci.* 4, 248–250. doi:10.1038/ngeo1095

622 Beyene, A., Abdelsalam, M.G., 2005. Tectonics of the Afar Depression: A review and synthesis.
 623 *J. African Earth Sci.* 41, 41–59. doi:10.1016/j.jafrearsci.2005.03.003

624 Bialas, R.W., Buck, W.R., Qin, R., 2010. How much magma is required to rift a continent? Earth
625 Planet. Sci. Lett. 292, 68–78. doi:10.1016/j.epsl.2010.01.021

626 Bindeman, I.N., Ponomareva, V. V., Bailey, J.C., Valley, J.W., 2004. Volcanic arc of
627 Kamchatka: a province with high- $\delta^{18}\text{O}$ magma sources and large-scale $^{18}\text{O}/^{16}\text{O}$ depletion of the
628 upper crust. *Geochim. Cosmochim. Acta* 68, 841–865. doi:10.1016/j.gca.2003.07.009

629 Bizouard, H., Barberi, F., Varet, J., 1980. Mineralogy and petrology of Erta Ale and Boina
630 volcanic series, Afar Rift, Ethiopia. *J. Petrol.* 21, 401–436. doi:10.1093/petrology/21.2.401

631 Bucholz, C.E., Jagoutz, O., VanTongeren, J.A., Setera, J., Wang, Z., 2017. Oxygen isotope
632 trajectories of crystallizing melts: Insights from modeling and the plutonic record. *Geochim.*
633 *Cosmochim. Acta* 207, 154–184. doi:10.1016/j.gca.2017.03.027

634 Buck, W.R., 2006. The role of magma in the development of the Afro-Arabian Rift System.
635 *Geol. Soc. London, Spec. Publ.* 259, 43–54. doi:10.1144/GSL.SP.2006.259.01.05

636 Cornwell, D.G., Mackenzie, G.D., England, R.W., Maguire, P.K.H., Asfaw, L.M., Oluma, B.,
637 2006. Northern Main Ethiopian Rift crustal structure from new high-precision gravity data. *Geol.*
638 *Soc. London, Spec. Publ.* doi:10.1144/GSL.SP.2006.259.01.23

639 d'Acremont, E., Leroy, S., Beslier, M.-O., Bellahsen, N., Fournier, M., Robin, C., Maia, M.,
640 Gente, P., 2005. Structure and evolution of the eastern Gulf of Aden conjugate margins from
641 seismic reflection data. *Geophys. J. Int.* 160, 869–890. doi:10.1111/j.1365-246X.2005.02524.x

642 Darling, W.G., Gizaw, B., Arusei, M.K., 1996. Lake-groundwater relationships and fluid-rock
643 interaction in the East African Rift Valley: isotopic evidence. *J. African Earth Sci.* 22, 423–431.
644 doi:10.1016/0899-5362(96)00026-7

645 Deering, C.D., Cole, J.W., Vogel, T.A., 2008. A rhyolite compositional continuum governed by
646 lower crustal source conditions in the taupo volcanic zone, New Zealand. *J. Petrol.* 49, 2245–
647 2276. doi:10.1093/petrology/egn067

648 Desissa, M., Johnson, N.E., Whaler, K.A., Hautot, S., Fisseha, S., Dawes, G.J.K., 2013. A
649 mantle magma reservoir beneath an incipient mid-ocean ridge in Afar, Ethiopia. *Nat. Geosci.* 6,
650 861–865. doi:10.1038/ngeo1925

651 Dufek, J., Bergantz, G.W., 2005. Lower crustal magma genesis and preservation: a stochastic
652 framework for the evaluation of basalt–crust interaction. *J. Petrol.* 46, 2167–2195.

653 Duffield, W.A., Bullen, T.D., Clynne, M.A., Fournier, R.O., Janik, C.J., Lanphere, M.A.,
654 Lowenstern, J., Smith, J.G., Giorgis, L.W., Kahsai, G., 1997. Geothermal potential of the Alid
655 volcanic center, Danakil Depression, Eritrea. US Geological Survey, Open File Report 97–291.

656 Ebinger, C., 2005. Continental break-up: the East African perspective. *Astron. Geophys.* 46,
657 2.16-2.21. doi:10.1111/j.1468-4004.2005.46216.x

658 Ebinger, C.J., Casey, M., 2001. Continental breakup in magmatic provinces: An Ethiopian
659 example. *Geology* 29, 527–530. doi:10.1130/0091-7613(2001)029<0527:CBIMPA>2.0.CO;2

660 Eiler, J.M., 2001. Oxygen Isotope Variations of Basaltic Lavas and Upper Mantle Rocks. *Rev.*
661 *Mineral. Geochemistry* 43, 319–364. doi:10.2138/gsrmg.43.1.319

662 Ferguson, D.J., Barnie, T.D., Pyle, D.M., Oppenheimer, C., Yirgu, G., Lewi, E., Kidane, T.,
663 Carn, S., Hamling, I., 2010. Recent rift-related volcanism in Afar, Ethiopia. *Earth Planet. Sci.*
664 *Lett.* 292, 409–418. doi:10.1016/j.epsl.2010.02.010

665 Ferguson, D.J., MacLennan, J., Bastow, I.D., Pyle, D.M., Jones, S.M., Keir, D., Blundy, J.D.,
666 Plank, T., Yirgu, G., 2013a. Melting during late-stage rifting in Afar is hot and deep. *Nature* 499,
667 70–3. doi:10.1038/nature12292

668 Ferguson, D.J., Calvert, A.T., Pyle, D.M., Blundy, J.D., Yirgu, G., Wright, T.J., 2013b.
669 Constraining timescales of focused magmatic accretion and extension in the Afar crust using
670 lava geochronology. *Nat. Commun.* 4, 1416. doi:10.1038/ncomms2410

671 Field, L., Blundy, J., Brooker, R. A., Wright, T., Yirgu, G., 2012a. Magma storage conditions
672 beneath Dabbahu Volcano (Ethiopia) constrained by petrology, seismicity and satellite geodesy.
673 *Bull. Volcanol.* doi:10.1007/s00445-012-0580-6

674 Field, L., Barnie, T., Blundy, J., Brooker, R.A., Keir, D., Lewi, E., Saunders, K., 2012b.
675 Integrated field, satellite and petrological observations of the November 2010 eruption of Erta
676 Ale. *Bull. Volcanol.* 74, 2251–2271. doi:10.1007/s00445-012-0660-7

677 Field, L., Blundy, J., Calvert, A., Yirgu, G., 2013. Magmatic history of Dabbahu, a composite
678 volcano in the Afar Rift, Ethiopia. *Bull. Geol. Soc. Am.* 125, 128–147. doi:10.1130/B30560.1

679 Furman, T., Bryce, J., Rooney, T.O., Hanan, B., Yirgu, G., Ayalew, D., 2006. Heads and tails: 30
680 million years of the Afar plume. *Geol. Soc. London, Spec. Publ.* 259, 95–119.

681 Gallacher, R.J., Keir, D., Harmon, N., Stuart, G., Leroy, S., Hammond, J.O.S., Kendall, J.-M.,
682 Ayele, A., Goitom, B., Ogubazghi, G., Ahmed, A., 2016. The initiation of segmented buoyancy-
683 driven melting during continental breakup. *Nat. Commun.* 7, 13110. doi:10.1038/ncomms13110

684 Giordano, F., D'Antonio, M., Civetta, L., Tonarini, S., Orsi, G., Ayalew, D., Yirgu, G.,
685 Dell'Erba, F., Di Vito, M. A., Isaia, R., 2014. Genesis and evolution of mafic and felsic magmas
686 at Quaternary volcanoes within the Main Ethiopian Rift: Insights from Gedemsa and Fanta 'Ale
687 complexes. *Lithos* 188, 130–144. doi:10.1016/j.lithos.2013.08.008

688 Gleeson, M.L.M., Stock, M.J., Pyle, D.M., Mather, T.A., Hutchison, W., Yirgu, G., Wade, J.,
689 2017. Constraining magma storage conditions at a restless volcano in the Main Ethiopian Rift
690 using phase equilibria models. *J. Volcanol. Geotherm. Res.* 337, 44–61.
691 doi:10.1016/j.jvolgeores.2017.02.026

692 Goodenough, K.M., Schilling, J., Jonsson, E., Kalvig, P., Charles, N., Tuduri, J., Deady, E.A.,
693 Sadeghi, M., Schiellerup, H., Müller, A., Bertrand, G., Arvanitidis, N., Eliopoulos, D.G., Shaw,
694 R.A., Thrane, K., Keulen, N., 2016. Europe's rare earth element resource potential: An overview
695 of REE metallogenic provinces and their geodynamic setting. *Ore Geol. Rev.* 72, 838–856.
696 doi:10.1016/j.oregeorev.2015.09.019

697 Graham, I.J., Cole, J.W., Briggs, R.M., Gamble, J.A., Smith, I.E.M., 1995. Petrology and
698 petrogenesis of volcanic rocks from the Taupo Volcanic Zone: a review. *J. Volcanol. Geotherm.*
699 *Res.* 68, 59–87. doi:10.1016/0377-0273(95)00008-I

700 Gualda, G.A.R., Ghiorso, M.S., Lemons, R. V., Carley, T.L., 2012. Rhyolite-MELTS: A
701 modified calibration of MELTS optimized for silica-rich, fluid-bearing magmatic systems. *J.*
702 *Petrol.* 53, 875–890. doi:10.1093/petrology/egr080

703 Hammond, J.O.S., Kendall, J.M., Stuart, G.W., Keir, D., Ebinger, C., Ayele, A., Belachew, M.,
704 2011. The nature of the crust beneath the Afar triple junction: Evidence from receiver functions.
705 *Geochemistry, Geophys. Geosystems* 12. doi:10.1029/2011GC003738

706 Harris, C., Smith, S.H., le Roex, A.P., 2000. Oxygen isotope composition of phenocrysts from
707 Tristan da Cunha and Gough Island lavas: variation with fractional crystallization and evidence
708 for assimilation. *Contrib. to Mineral. Petrol.* 138, 164–175. doi:10.1007/s004100050015

709 Hayward, N.J., Ebinger, C.J., 1996. Variations in the along-axis segmentation of the Afar Rift
710 system. *Tectonics* 15, 244. doi:10.1029/95TC02292

711 Hofmann, C., Courtillot, V., Féraud, G., Rochette, P., Yirgu, G., Ketefo, E., Pik, R., 1997.
712 Timing of the Ethiopian flood basalt event and implications for plume birth and global change.
713 *Nature* 389, 838–841. doi:10.1038/39853

714 Hutchison, W., Fusillo, R., Pyle, D.M., Mather, T.A., Blundy, J.D., Biggs, J., Yirgu, G., Cohen,
715 B.E., Brooker, R.A., Barfod, D.N., Calvert, A.T., 2016a. A pulse of mid-Pleistocene rift
716 volcanism in Ethiopia at the dawn of modern humans. *Nat. Commun.* 7, 13192.
717 doi:10.1038/ncomms13192

718 Hutchison, W., Biggs, J., Mather, T.A., Pyle, D.M., Lewi, E., Yirgu, G., Caliro, S., Chiodini, G.,
719 Clor, L.E., Fischer, T.P., 2016b. Causes of unrest at silicic calderas in the East African Rift: New

720 constraints from InSAR and soil-gas chemistry at Aluto volcano, Ethiopia. *Geochemistry,*
721 *Geophys. Geosystems* 17, 3008–3030. doi:10.1002/2016GC006395

722 Hutchison, W., Pyle, D.M., Mather, T.A., Yirgu, G., Biggs, J., Cohen, B.E., Barfod, D.N., Lewi,
723 E., 2016c. The eruptive history and magmatic evolution of Aluto volcano: new insights into
724 silicic peralkaline volcanism in the Ethiopian rift. *J. Volcanol. Geotherm. Res.* 328, 9–33.
725 doi:10.1016/j.jvolgeores.2016.09.010

726 Irvine, T.N., Baragar, W.R.A., 1971. A Guide to the Chemical Classification of the Common
727 Volcanic Rocks. *Can. J. Earth Sci.* 8, 523–548. doi:10.1139/e71-055

728 Karakas, O., Dufek, J., 2015. Melt evolution and residence in extending crust: Thermal modeling
729 of the crust and crustal magmas. *Earth Planet. Sci. Lett.* 425, 131–144.
730 doi:10.1016/j.epsl.2015.06.001

731 Keir, D., Bastow, I.D., Corti, G., Mazzarini, F., Rooney, T.O., 2015. The origin of along-rift
732 variations in faulting and magmatism in the Ethiopian Rift. *Tectonics* 34, 464–477.
733 doi:10.1002/2014TC003698

734 Keranen, K., Klemperer, S.L., Gloaguen, R., EAGLE Working Group, 2004. Three-dimensional
735 seismic imaging of a protoridge axis in the Main Ethiopian rift. *Geology* 32, 949.
736 doi:10.1130/G20737.1

737 Lahitte, P., Gillot, P.-Y., Kidane, T., Courtillot, V., Bekele, A., 2003. New age constraints on the
738 timing of volcanism in central Afar, in the presence of propagating rifts. *J. Geophys. Res. Solid*
739 *Earth* 108. doi:10.1029/2001JB001689

740 Lewi, E., Keir, D., Birhanu, Y., Blundy, J., Stuart, G., Wright, T., Calais, E., 2016. Use of a
741 high-precision gravity survey to understand the formation of oceanic crust and the role of melt at
742 the southern Red Sea rift in Afar, Ethiopia. *Geol. Soc. London, Spec. Publ.* 420, 165–180.
743 doi:10.1144/SP420.13

744 Macaulay, C.I., Fallick, A. E., Haszeldine, R.S., Graham, C.M., 2000. Methods of laser-based
745 stable isotope measurement applied to diagenetic cements and hydrocarbon reservoir quality.
746 *Clay Miner.* 35, 313–313. doi:10.1180/000985500546684

747 Maccaferri, F., Rivalta, E., Keir, D., Acocella, V., 2014. Off-rift volcanism in rift zones
748 determined by crustal unloading. *Nat. Geosci.* 7, 297–300. doi:10.1038/ngeo2110

749 Macdonald, R., 1974. Nomenclature and petrochemistry of the peralkaline oversaturated
750 extrusive rocks. *Bull. Volcanol.* 38, 498–516. doi:10.1007/BF02596896

751 Macdonald, R., 2012. Evolution of peralkaline silicic complexes: Lessons from the extrusive
752 rocks. *Lithos* 152, 11–22. doi:10.1016/j.lithos.2012.01.014

753 Macdonald, R., Davies, G.R., Bliss, C.M., Leat, P.T., Bailey, D.K., Smith, R.L., 1987.
754 Geochemistry of high-silica peralkaline rhyolites, Naivasha, Kenya rift valley. *J. Petrol.* 28, 979–
755 1008. doi:10.1093/petrology/28.6.979

756 Macdonald, R., Belkin, H.E., Fitton, J.G., Rogers, N.W., Nejbert, K., Tindle, A.G., Marshall,
757 A.S., 2008. The roles of fractional crystallization, magma mixing, crystal mush remobilization
758 and volatile-melt interactions in the genesis of a young basalt-peralkaline rhyolite suite, the

759 greater Olkaria volcanic complex, Kenya Rift valley. *J. Petrol.* 49, 1515–1547.
 760 doi:10.1093/petrology/egn036

761 Macdonald, R., Bagiński, B., Upton, B.G.J., 2014. The volcano-pluton interface; The Longonot
 762 (Kenya) and Kûngnât (Greenland) peralkaline complexes. *Lithos* 196–197, 232–241.
 763 doi:10.1016/j.lithos.2014.03.009

764 Mackenzie, G.D., Thybo, H., Maguire, P.K.H., 2005. Crustal velocity structure across the Main
 765 Ethiopian Rift: results from two-dimensional wide-angle seismic modelling. *Geophys. J. Int.*
 766 162, 994–1006. doi:10.1111/j.1365-246X.2005.02710.x

767 Maguire, P.K.H., Keller, G.R., Klemperer, S.L., Mackenzie, G.D., Keranen, K., Harder, S.,
 768 O'Reilly, B., Thybo, H., Asfaw, L., Khan, M.A., Amha, M., 2006. Crustal structure of the
 769 northern Main Ethiopian Rift from the EAGLE controlled-source survey; a snapshot of incipient
 770 lithospheric break-up. *Geol. Soc. London, Spec. Publ.* 259, 269–292.
 771 doi:10.1144/GSL.SP.2006.259.01.21

772 Mahatsente, R., Jentzsch, G., Jahr, T., 1999. Crustal structure of the Main Ethiopian Rift from
 773 gravity data: 3-dimensional modeling. *Tectonophysics* 313, 363–382. doi:10.1016/S0040-
 774 1951(99)00213-9

775 Makris, J., Ginzburg, A., 1987. The Afar Depression: transition between continental rifting and
 776 sea-floor spreading. *Tectonophysics* 141, 199–214. doi:10.1016/0040-1951(87)90186-7

777 Marshall, A.S., MacDonald, R., Rogers, N.W., Fitton, J.G., Tindle, A.G., Nejbert, K., Hinton,
 778 R.W., 2009. Fractionation of peralkaline silicic magmas: The greater olkaria volcanic complex,
 779 Kenya Rift Valley. *J. Petrol.* 50, 323–359. doi:10.1093/petrology/egp001

780 Martin, E., Sigmarsson, O., 2010. Thirteen million years of silicic magma production in Iceland:
 781 Links between petrogenesis and tectonic settings. *Lithos* 116, 129–144.
 782 doi:10.1016/j.lithos.2010.01.005

783 Medynski, S., Pik, R., Burnard, P., Vye-Brown, C., France, L., Schimmelpfennig, I., Whaler, K.,
 784 Johnson, N., Benedetti, L., Ayelew, D., Yirgu, G., 2015. Stability of rift axis magma reservoirs:
 785 Spatial and temporal evolution of magma supply in the Dabbahu rift segment (Afar, Ethiopia)
 786 over the past 30 kyr. *Earth Planet. Sci. Lett.* 409, 278–289. doi:10.1016/j.epsl.2014.11.002

787 Medynski, S., Pik, R., Burnard, P., Williams, a., Vye-Brown, C., Ferguson, D., Blard, P.H.,
 788 France, L., Yirgu, G., Seid, J.I., Ayalew, D., Calvert, A., 2013. Controls on magmatic cycles and
 789 development of rift topography of the Manda Hararo segment (Afar, Ethiopia): Insights from
 790 cosmogenic ³He investigation of landscape evolution. *Earth Planet. Sci. Lett.* 367, 133–145.
 791 doi:10.1016/j.epsl.2013.02.006

792 Norton, D., Taylor, H.P., 1979. Quantitative simulation of the hydrothermal systems of
 793 crystallizing magmas on the basis of transport theory and oxygen isotope data: An analysis of the
 794 Skaergaard intrusion. *J. Petrol.* 20, 421–486. doi:10.1093/petrology/20.3.421

795 Pagli, C., Wright, T.J., Ebinger, C.J., Yun, S.-H., Cann, J.R., Barnie, T., Ayele, A., 2012.
 796 Shallow axial magma chamber at the slow-spreading Erta Ale Ridge. *Nat. Geosci.* 5, 284–288.
 797 doi:10.1038/ngeo1414

798 Peccerillo, A., Barberio, M.R., Yirgu, G., Ayalew, D., Barbieri, M., Wu, T.W., 2003.
799 Relationships between Mafic and Peralkaline Silicic Magmatism in Continental Rift Settings: a
800 Petrological, Geochemical and Isotopic Study of the Gedemsa Volcano, Central Ethiopian Rift.
801 J. Petrol. 44, 2003–2032. doi:10.1093/petrology/egg068

802 Peccerillo, A., Donati, C., Santo, A., Orlando, A., 2007. Petrogenesis of silicic peralkaline rocks
803 in the Ethiopian rift: geochemical evidence and volcanological implications. J. African Earth 48,
804 161–173. doi:10.1016/j.jafrearsci.2006.06.010

805 Peccerillo, A., Mandefro, B., Solomon, G., Bedru, H., Tesfaye, K., 1998. The Precambrian rocks
806 from Southern Ethiopia: petrology, geochemistry and their interaction with the recent volcanism
807 from the Ethiopian Rift Valley. Neues Jahrb. für Mineral. 237–262.

808 Pope, E.C., Bird, D.K., Arnórsson, S., 2013. Evolution of low- ^{18}O Icelandic crust. Earth Planet.
809 Sci. Lett. 374, 47–59. doi:10.1016/j.epsl.2013.04.043

810 Rampey, M.L., 2005. The geology of the Kone volcanic complex, Ethiopia. University of
811 Cambridge, PhD Thesis.

812 Rampey, M.L., Oppenheimer, C., Pyle, D.M., Yirgu, G., 2014. Journal of African Earth Sciences
813 Physical volcanology of the Gubisa Formation, Kone Volcanic Complex, Ethiopia. J. African
814 Earth Sci. 96, 212–219. doi:10.1016/j.jafrearsci.2014.04.009

815 Rampey, M.L., Oppenheimer, C., Pyle, D.M., Yirgu, G., 2010. Caldera-forming eruptions of the
816 Quaternary Kone Volcanic Complex, Ethiopia. J. African Earth Sci. 58, 51–66.
817 doi:10.1016/j.jafrearsci.2010.01.008

818 Rooney, T., Furman, T., Bastow, I., Ayalew, D., Yirgu, G., 2007. Lithospheric modification
 819 during crustal extension in the Main Ethiopian Rift. *J. Geophys. Res.* 112.
 820 doi:10.1029/2006JB004916

821 Rooney, T.O., 2017. The Cenozoic magmatism of East-Africa: Part I — Flood basalts and pulsed
 822 magmatism. *Lithos* 286–287, 264–301. doi:10.1016/j.lithos.2017.05.014

823 Rooney, T.O., Herzberg, C., Bastow, I.D., 2012a. Elevated mantle temperature beneath East
 824 Africa. *Geology* 40, 27–30. doi:10.1130/G32382.1

825 Rooney, T.O., Hart, W.K., Hall, C.M., Ayalew, D., Ghiorso, M.S., Hidalgo, P., Yirgu, G., 2012b.
 826 Peralkaline magma evolution and the tephra record in the Ethiopian Rift. *Contrib. to Mineral.*
 827 *Petrol.* 164, 407–426. doi:10.1007/s00410-012-0744-6

828 Rooney, T.O., Hanan, B.B., Graham, D.W., Furman, T., Blichert-toft, J., Schilling, J.G., 2012c.
 829 Upper mantle pollution during Afar plume-continental rift interaction. *J. Petrol.* 53, 365–389.
 830 doi:10.1093/petrology/egr065

831 Saria, E., Calais, E., Stamps, D.S., Delvaux, D., Hartnady, C.J.H., 2014. Present-day kinematics
 832 of the East African Rift. *J. Geophys. Res. Solid Earth* 119, 3584–3600.
 833 doi:10.1002/2013JB010901

834 Scaillet, B., Macdonald, R., 2001. Phase relations of peralkaline silicic magmas and petrogenetic
 835 implications. *J. Petrol.* 42, 825–845. doi:10.1093/petrology/42.4.825

836 Sharp, Z., 1990. A laser-based microanalytical method for the in situ determination of oxygen
837 isotope ratios of silicates and oxides. *Geochim. Cosmochim. Acta* 54, 1353–1357.
838 doi:10.1016/0016-7037(90)90160-M

839 Teklay, M., Kröner, A., Mezger, K., Oberhänsli, R., 1998. Geochemistry, Pb-Pb single zircon
840 ages and Nd-Sr isotope composition of Precambrian rocks from southern and eastern Ethiopia:
841 implications for crustal evolution in East Africa. *J. African Earth Sci.* 26, 207–227.
842 doi:10.1016/S0899-5362(98)00006-2

843 Teklemariam, M., Battaglia, S., Gianelli, G., Ruggieri, G., 1996. Hydrothermal alteration in the
844 Aluto-Langano geothermal field, Ethiopia. *Geothermics* 25, 679–702. doi:10.1016/S0375-
845 6505(96)00019-3

846 Watson, E.B., 1979. Zircon saturation in felsic liquids: Experimental results and applications to
847 trace element geochemistry. *Contrib. to Mineral. Petrol.* 419, 407–419.

848 White, J.C., Parker, D.F., Ren, M., 2009. The origin of trachyte and pantellerite from Pantelleria,
849 Italy: Insights from major element, trace element, and thermodynamic modelling. *J. Volcanol.*
850 *Geotherm. Res.* 179, 33–55. doi:10.1016/j.jvolgeores.2008.10.007

851 Wolfenden, E., Ebinger, C., Yirgu, G., Deino, A., Ayalew, D., 2004. Evolution of the northern
852 Main Ethiopian rift: birth of a triple junction. *Earth Planet. Sci. Lett.* 224, 213–228.
853 doi:10.1016/j.epsl.2004.04.022

854 Wolfenden, E., Ebinger, C., Yirgu, G., Renne, P.R., Kelley, S.P., 2005. Evolution of a volcanic
855 rifted margin: Southern Red Sea, Ethiopia. *Geol. Soc. Am. Bull.* 117, 846. doi:10.1130/B25516.1

Xu, W., Rivalta, E., Li, X., 2017. Magmatic architecture within a rift segment: Articulate axial magma storage at Erta Ale volcano, Ethiopia. *Earth Planet. Sci. Lett.* 476, 79–86.
doi:10.1016/j.epsl.2017.07.051

Figure Captions

Figure 1. Topographic maps summarizing geotectonic setting and crustal structure in Ethiopia. In (a) the black lines represent major plate boundaries, coloured symbols identify the volcanic systems considered in this study, and the grey square corresponds to the inset b. The major terrestrial rift zones are: the Afar Rift (AR); the Main Ethiopian Rift (MER) and Kenyan Rift (KR). Within each rift volcanic activity is localized in magmatic segments. Dabbahu and Badi are located on and off the Manda Hararo Rift Segment (MHRS) of the Afar Rift (inset b), while the Erta Ale range lies in the Erta Ale Segment (EAS). Aluto, Gedemsa and Kone are located along the MER, but Kone lies in a different magmatic segment from Aluto and Gedemsa. White arrows show current extension vectors relative to a fixed Nubian Plate (after Saria et al., 2014). (b) Hillshade digital elevation model of the MHRS showing the location of Dabbahu and Badi silicic volcanic systems. The black dashed line identifies the rift axis and the orange area represents the focus of recent dyke intrusion, faulting and eruptions (Ferguson et al., 2010). (c) A summary of crustal sections adjacent to the volcanic systems considered in this study. Sections are based on controlled source seismic experiments and receiver function studies (after Hammond et al., 2011 and references therein). Section locations are shown as circled numbers on the topographic map (a). Approximate depths of silicic magma reservoirs are based on

deformation (Hutchison et al., 2016c), seismic and petrological constraints (Field et al., 2012a; Gleeson et al., 2017).

Figure 2. Selected major element classification diagrams (all data are anhydrous normalized).

(a) Total alkalis versus silica (TAS) diagram. The grey dashed line shows the alkaline sub-alkaline (tholeiitic) divide of Irvine and Baragar (1971). Rhyolites with low alkalis (<8 wt. %) often have anomalous low Na₂O and high loss on ignition values suggestive of post-emplacment alteration (Peccerillo et al., 2003). (b) Peralkalinity index (PI) versus Aluminium Saturation Index (ASI) is used to define metaluminous, peraluminous and peralkaline composition fields for silicic rocks. PI is defined as the molar (Na₂O+K₂O) / Al₂O₃ while ASI is defined as the molar ratio Al₂O₃ / (CaO + K₂O + Na₂O). For each volcanic system the most peralkaline samples are labelled. (c) Peralkaline classification diagram (Al₂O₃ versus FeO_t) based on Macdonald (1974).

Figure 3. Major element Harker diagrams of whole-rock compositions determined by XRF.

Best-fit Rhyolite-MELTS liquid lines of descent for Aluto and Dabbahu are overlain (black dashed lines). The model parameters are provided in Table 3. Inaccuracies in apatite crystallization for peralkaline systems in Rhyolite-MELTS explain the poor fit between the data and models for P₂O₅. A schematic showing a more likely liquid line of descent for P₂O₅ is provided and illustrates how mixed samples fall beneath the peak. Natural samples that show textural evidence for magma mixing are circled by the grey dashed line.

Figure 4. Selected trace element compositions for the volcanic systems. The influence of

magmatic differentiation and certain fractionating mineral phases are shown by the black arrows.

Data for ~30 Ma comendite ignimbrites associated with flood basalt volcanism (after, Ayalew et

al., 2002 and Ayalew and Yirgu, 2003) are shown as dark grey shaded fields (with asterisk in bottom right hand plot). Pan-African Precambrian crust trace element data are from Peccerillo et al. (1998).

Figure 5. Isotopic constraints on magma petrogenesis. **a)** $\delta^{18}\text{O}_{\text{melt}}$ evolution predicted by model fractional crystallization (FC) scenarios compared to data from across Ethiopia. For clarity samples are arranged according to their rock type (total alkalis-silica classification) so that fractionation increases from left to right. Data point colours relate to each volcanic system; symbol style corresponds to the material analysed. To correct $\delta^{18}\text{O}_{\text{mineral}}$ to $\delta^{18}\text{O}_{\text{melt}}$ we used experimentally determined fraction factors of Appora et al. (2003). The $\delta^{18}\text{O}_{\text{melt}}$ trajectory for our best-fit Rhyolite-MELTS models is shown by the grey shaded region (labelled FC trajectory). Fractional crystallization scenarios predict a moderate increase in $\delta^{18}\text{O}_{\text{melt}}$ assuming a parental melt range of 5.0–5.8 ‰ (see text for detailed discussion). Samples that fall off the FC trajectory have been contaminated by Pan-African Precambrian crustal rocks (high $\delta^{18}\text{O}$) or hydrothermally altered crust (low $\delta^{18}\text{O}$), which we have assigned end-member values of 12 ‰ and 2 ‰ respectively (Section 5.1). Orange and blue lines above and below the FC trajectory provide estimates for the amount of end-member crustal melt that would need to be assimilated in order to explain $\delta^{18}\text{O}_{\text{melt}}$. New $\delta^{18}\text{O}$ are presented in Table 2. Erta Ale and MHRS $\delta^{18}\text{O}$ are from Barrat et al. (1998, 2003). ~30 Ma ignimbrites are from Ayalew et al. (2002). **b)** ϵ_{Nd} and $^{87}\text{Sr}/^{86}\text{Sr}$ constraints on crustal assimilation in Ethiopia. Symbol colours are the same as a), while symbol style corresponds to rock type. All isotopes for ~30 Ma samples are age-corrected (see Rooney, 2017, and references therein). New analyses are from Dabbahu, Badi and MHRS while previous analyses are compiled from: Barrat et al. (1998, 2003); Peccerillo et al., (2003, 2007); Furman et al. (2006); Giordano et al. (2014) and Ayalew et al., (2016). The field of MER and Afar Rift

Quaternary basalts (assumed representative of Ethiopian mantle) are outlined by the black dotted line (see Ayalew et al., 2016 and references therein). A Pan-African lithosphere endmember (after Rooney et al., 2012c) is shown by the grey star while a typical uncontaminated mantle-derived melt from Afar is shown by the white star. Binary mixing models for pristine mantle-derived pantellerite and basalt compositions with Pan-African crust are shown by the black lines at increments of 5 %. Mixing of partial melts of granitoid rocks (equivalent to Pan-African crustal materials using batch melting models) with uncontaminated-endmembers produced similar trends to those shown and for figure clarity these are not displayed. The end-members used in the model had Sr and Nd concentrations as follows: pantellerite: 5 and 200 ppm; basalt: 500 and 30 ppm and Pan-African: 225 and 22 ppm (in accordance with Rooney et al., 2012c).

Figure 6. Comparison of erupted volumes, fluxes and melt evolution for the volcanic systems and a summary of key findings. **a)** The total erupted volume of each volcanic system (Table 4) is shown by the upper coloured bars and is plotted against local crustal thickness (after Hammond et al., 2011). Cumulate volumes are shown by the lower, darker bars and were calculated using Rhyolite-MELTS models (see text). **b)** Eruptive and cumulate fluxes are shown for each system and were calculated using available age constraints (Tables 1, 4). Two flux bars are shown for Dabbahu, using two estimates for the age of the system (300 and 500 ka, Tables 1, 4). **c)** The average Zr content of the most evolved samples for each volcanic system plotted against the erupted volume flux (b). Each system is represented by a different coloured symbol. To get a representative sample of the most evolved components of each volcanic system we average the Zr concentration of the silicic samples with highest Zr (which we define as the upper quartile of the data). There is a clear negative relationship between eruptive flux and average Zr (i.e. melt evolution). This suggests that local magma supply is an important control on fractional

945 crystallization and melt evolution processes. **d)** Schematic summarizing crustal structures of the
946 EAS, MHRS and MER and petrological constraints on magmagenesis. Grey colours represent
947 intrusive melts while orange rectangles represent cumulate bodies. Blue spirals reflect
948 hydrothermal circulation that generates low $\delta^{18}\text{O}$ crust. Abbreviations: RF: rift fill (i.e., lavas and
949 sedimentary cover); UC: upper crust; LC: lower crust. Intermediate-mature continental rift zones
950 (e.g., MER and MHRS) promote extreme melt evolution since parental basalts are transitional in
951 composition and local magmatic flux is low.

952

Table 1. Summary of the geological setting, eruptive history and age constraints for the six volcanic systems considered in this study. Note the following abbreviations: MER: Main Ethiopian Rift; AG: Aluto-Gedemsa; BK: Boseti-Kone; MHRS: Manda Hararo Rift Segment and EAS: Erta Ale Segment.

rift zone	volcanic system (magmatic segment)	eruptive history/setting	age constraints	key references
MER	Aluto (AG segment)	Initially built up as a trachytic complex before undergoing caldera collapse. Significant volumes of post-caldera silicic volcanism then took place rebuilding the main edifice.	Caldera forming eruption took place at 316 ± 19 ka and 306 ± 12 ka; post-caldera volcanism initiated at 55 ± 19 ka with recent eruptions in last 1 ka	Hutchison et al., 2016a, b; Gleeson et al., 2017
	Gedemsa (AG segment)	Initial activity built up a trachytic-rhyolitic complex. Caldera collapse then took place possibly via multiple explosive eruptions. Finally, a set of small coalescing post-caldera silicic edifices and fissure basalts were erupted in the centre of the caldera.	K/Ar dating places pre-caldera samples 320 ± 20 ka and post-caldera samples 260 ± 20 ka	Peccerillo et al., 2003; Giordano et al., 2014
	Kone (BK segment)	Pre-caldera activity built up a trachytic cone. Then four major eruptive phases took place and generated a large collapse scar. Post-caldera mafic eruptions have partially infilled the collapse and very minor silicic eruptions have occurred around the ring fracture.	None	Rampey et al. 2010, 2014
Afar Rift	Badi (MHRS)	Badi is an off-axis silicic volcano (Figure 1b) - it is composed of coalescing lava flows and domes and no caldera collapse scars are observed.	Earliest eruptions dated at 290 ± 4 ka (K/Ar age for a rhyolitic flow near base of edifice), recent basaltic lavas dated between ~140 and 25 ka	Lahitte et al., 2003; Ferguson et al., 2013b
	Dabbahu (MHRS)	Dabbahu is an on-axis volcano (Figure 1b). The central edifice comprises coalescing flows and domes that span an almost complete basalt-peralkaline rhyolite compositional series. No large-volume ignimbrite deposits or caldera features have yet been identified.	$^{40}\text{Ar}/^{39}\text{Ar}$ indicate multiple eruptive cycles between 60 and 5 ka, cosmogenic ^3He ages suggest Dabbahu has been active for >100 ka	Barberi et al., 1975; Field et al., 2012a, 2013; Medynski et al., 2013
	Erta Ale (EAS)	The ~80 km Erta Ale range forms an axial volcanic ridge comprising six distinct eruptive centres (Gada Ale, Alu-Dalafilla, Borale Ale, Erta Ale, Hayli Gubbi and Ale Bagu). Magma compositions range from transitional alkali-tholeiitic basalt to rhyolite.	Age constraints on onset of volcanism are limited, oldest K/Ar are ~1000 ka	Barberi and Varet, 1970; Barberi et al., 1972; Bizourd et al., 1980; Barrat et al., 1998; Beyene and Abdelsalam, 2005

958 **Table 2.** New oxygen isotope data for Ethiopian volcanic systems.

volcanic system ^a	sample code	rock type ^b	material analysed	$\delta^{18}\text{O}$ (‰ V-SMOW)	age (stratigraphic context) ^c
Aluto	15-02-09	trachyte	glass	6.5	570–330 ka (pre-caldera) ¹
	01-02-14	pantellerite	glass	6.2	62 ± 13 ka (post-caldera) ¹
	18-02-04	comendite	glass	6.1	55 ± 19 ka (post-caldera) ¹
	02-02-12	pantellerite	glass	6.2	60–10 ka (post-caldera) ¹
	31-01-LE	pantellerite	glass	6.4	16 ± 14 ka (post-caldera) ¹
	15-01-07B	TA	glass	5.2	n.d. (post-caldera) ¹
	18-11-01	BTA	glass	6.0	n.d. (post-caldera) ¹
	01-02-13	pantellerite	glass	6.2	<10 ka (post-caldera) ¹
Gedemsa ^d	GD22	rhyolite	alkali feldspar	7.9	>300 ka (pre-caldera) ²
	GD3	basalt	olivine	5.8	<300 ka (post-caldera) ²
	GD8	pantellerite	alkali feldspar	6.2	<300 ka (post-caldera) ²
	GD12	pantellerite	alkali feldspar	7.3	<300 ka (post-caldera) ²
	GD15	salic xenolith	alkali feldspar	6.8	n.d. (enclave in post-caldera rocks) ²
Kone	2-11	rhyolite	glass	7.1	n.d. (pre-caldera) ³
	3-30	pantellerite	glass	5.9	n.d. (post-caldera) ³
	4-11	pantellerite	glass	6.2	n.d. (post-caldera) ³
	3-34	pantellerite	glass	6.1	n.d. (post-caldera) ³
	4-21	basalt	glass	5.2	n.d. (post-caldera) ³
	2-20	basalt	glass	5.5	n.d. (post-caldera) ³
	3-31	basalt	glass	5.6	n.d. (post-caldera) ³
	2-22	basalt	glass	5.4	n.d. (post-caldera) ³
Badi	3016	basalt	glass	6.1	n.d. ⁴
	29 09	TA	glass	6.0	138.9 ± 1.1 ka ⁴
	29 11	basalt	glass	5.3	54.5 ± 5.6 ka ⁴
	Bad 4	basalt	olivine	4.8	39.8 ± 3.7 ka ⁵
	Bad 5	basalt	olivine	5.0	39.8 ± 3.7 ka ⁵
	BADI 05	basalt	glass	5.1	25.4 ± 9.4 ka ⁴
	24 02	pantellerite	glass	5.6	n.d. ⁴
	30 12	pantellerite	glass	6.0	n.d. ⁴
	03 04	pantellerite	glass	6.5	n.d. ⁴
	BADI 07	pantellerite	glass	6.2	n.d. ⁴
MHRS axis	Dik 2	basalt	olivine	5.2	68.2 ± 3.3 ka ⁵
	Dik 3	basalt	olivine	5.2	68.2 ± 3.3 ka ⁵
	Gab I	basalt	olivine	5.1	66.1 ± 5.4 ka ⁵
	Gab G3	basalt	olivine	5.2	<2 ka ⁵
	2007	basalt	glass	5.3	2007 (fissure eruption) ⁶
	Gab B	basalt	olivine	5.3	n.d. ⁵
Dabbahu	LFAF 044	basalt	glass	5.4	>60 ka ⁷
	LFAF 045	basalt	glass	4.9	>60 ka ⁷
	LFAF 028	TB	glass	5.3	63–48 ka ⁷
	LFAF 106	TA	glass	6.4	63–48 ka ⁷
	LFAF 047a	BTA	glass	6.0	~49 ka ⁷
	LFAF 054	commendite	glass	5.7	30.1 ± 0.4 ka ⁷
	LFAF 011	comendite	glass	7.8	28.6 ± 0.7 ka ⁷
	LFAF 055	pantellerite	glass	5.7	<10 ka ⁷
	LFAF 063	pantellerite	glass	5.5	<10 ka ⁷

959 ^a Erta Ale $\delta^{18}\text{O}$ data (not shown) was presented in Barrat et al. (1998)

960 ^b Rock type abbreviations include: TB: trachybasalt; BTA: basaltic trachyandesite and TA: trachyandesite.

961 ^c Age and stratigraphic constraints are provided (when available) in the right-hand column. n.d. indicates that
962 no age data is currently available. Superscripts refer to prior published samples: 1: Hutchison et al. (2016a, b);
963 2: Peccerillo et al. (2003); 3: Rampey (2005); 4: Ferguson et al. (2013b); 5: Medynski et al. (2013); 6:
964 Ferguson et al. (2010) and 7: Field et al. (2013).

965 ^d Gedemsa $\delta^{18}\text{O}$ data were generated by Peccerillo et al. (2003). Note that sample labels for $\delta^{18}\text{O}$ have been
966 updated here following discussion with the authors (Peccerillo pers. comm.).

967

Table 3. Best-fit magma storage conditions for Aluto and Dabbahu constrained via Rhyolite-MELTS models and the range of input parameters that were explored. Note that we use a minimization routine developed by Gleeson et al. (2017) to identify the model storage conditions that most closely match the whole-rock compositions of the pantellerites (i.e., the most chemically-evolved samples).

		Aluto	Dabbahu
parameter range explored	pressure (MPa)	50–300	50–400
	H ₂ O content (wt. %)	0.5–3	0.5–2
	f _{O₂} (log units relative to QFM)	–2 to +1	0 to –1
Rhyolite-MELTS best-fit	pressure (MPa)	150	150
	H ₂ O content (wt. %)	0.5	0.5
	f _{O₂} (log units relative to QFM)	0	0
	starting composition	primitive rift-related basalt (17-01-05) sampled north-east of the complex	high MgO (>8 wt. %) basaltic sample LFAF 044 (Field et al., 2013)
	bulk density of crystal cumulate (kg m ^{–3})	3060	3040
petrological constraints		granophyric textures in some rhyolitic samples indicate crystallisation at shallow depths (~100 MPa)	melt inclusions suggest shallow storage <250 MPa. Fe-Ti oxides constrain f _{O₂} between QFM and QFM-1. 4±1 wt. % H ₂ O in rhyolites suggests ~0.5% H ₂ O in parent
relevant studies		Hutchison et al., 2016; Gleeson et al., 2017	Field et al., 2012a, 2013

Table 4. Erupted and cumulate volume estimates for the volcanic systems considered in this study.

volcanic system (eruption onset estimate) ^a	erupted volumes (from DEM) ^b		cumulate volumes (from Rhyolite- MELTS) ^c		cumulate volume: erupted volume	local rift volume filled by cumulate (%) ^d
	deposit volume (km ³)	eruptive flux (km ³ ka ⁻¹)	cumulate volume (km ³)	cumulate flux (km ³ ka ⁻¹)		
Aluto (500 ka)	pre-caldera edifice	48	145			
	caldera collapse	20	60			
	post-caldera	32	98			
	total	99	302	0.60	3.0	23
Gedemsa (500 ka)	pre-caldera edifice	33	135			
	caldera collapse	30	122			
	post-caldera	1	6			
	total	64	263	0.53	4.1	30
Kone (500 ka)	pre-caldera edifice	10	54			
	caldera collapse	15	75			
	total	25	129	0.26	5.2	25
Badi (300 ka)	edifice	29	92	0.31	3.2	9
Dabbahu (500–300 ka)	edifice	114	454	0.91–1.51	4.0	16–27
Erta Ale (1000 ka)	magmatic segment	405	209	0.21	0.5	1

^a The timing of eruption onsets at each system are not known precisely; we used best-estimates from existing geochronologies (Table 1). For the MER volcanoes, age constraints suggest initial eruptions at ~500 ka (i.e., between 750 and 300 ka, Hutchison et al., 2016a). At Badi we use a K/Ar age of ~300 ka from the base as the eruption onset (Lahitte et al., 2003). At Dabbahu, the oldest deposits are poorly exposed and although cosmogenic ³He ages (Medynski et al., 2013) indicate ages >100 ka no precise age is known for eruption onset. Instead we evaluate a range of values between 300 ka (assuming Dabbahu is equivalent in age to neighbouring Badi) and 500 ka (i.e. the minimum age of basalts on external margins of the MHRS which Dabbahu overlaps, Lahitte et al., 2003). The onset of volcanism in the Erta Ale magmatic segment is generally placed at less than ~1000 ka (see review of Beyene and Abdelsalam, 2005, based on K/Ar dating from Barberi et al., 1972).

^b Erupted volumes are calculated from digital elevation models (DEMs) following methods outlined by Hutchison et al., (2016a). At Aluto and Gedemsa it is possible to identify pre-, syn- and post-caldera phases. At Kone, post-caldera deposits are widely distributed but very minor in areal extent and have not been included in the long-term eruptive volume budget. At Badi and Dabbahu, caldera collapse phases are poorly

992 documented so we calculate the volume of the edifice. At Erta Ale, we calculated the volume of the entire
993 magmatic segment, which includes six volcanic centres: Gada Ale, Alu-Dalafilla, Borale Ale, Erta Ale, Hayli
994 Gubbi and Ale Bagu. We estimate cumulate volumes from whole rock Zr data, and Rhyolite-MELTS models
995 (details in Section 5.4).

996 ^c Cumulate fluxes and volumes are minima because they are based on erupted volumes and exclude any
997 evolved magmas that did not erupt.

998 ^d Percentage of rift volume (created by extension) that would be filled by magmatic cumulate. To estimate the
999 volume created by extension at each volcanic system we use the local crustal thickness (Figure 6a), extension
1000 rate (Figure 2) and their along-axis length (measured from DEMs).











